Soybean leaf physiological responses to drought stress improved via enhanced seed zinc and iron concentrations

Mohsen Movahhedi Dehnavi^{*} and Marzieh Jalil Sheshbahre Department Agronomy and Plant Breeding, Faculty of Agriculture, Yasouj University, Yasouj, (Received: 14/10/2015-Accepted: 27/01/2016)

Abstract:

To study the effects of both drought stress and seed zinc and iron concentrations on the photosynthesis, chlorophyll fluorescence, and proline and carbohydrate accumulations in soybean (*Glycine max* (L.) Merr. cv. M9) leaf, a split plot experiment was carried out in 2012 based on a randomized complete block design with three replications in greenhouse conditions at Yasouj University. Drought stress, based on 70% depletion of the available soil water, was designated as the main-plot that included drought stress at the flowering, pod filling, and seed filling stages. Zinc and iron concentrations were designated as sub-plot factors that included high zinc, iron and zinc+iron contents of the seed, and control as well as seed soaking in distilled water, 3% zinc sulfate, 3% iron sulfate, and 3% zinc+3% iron sulfate solutions. Results showed that photosynthesis characteristics, net photosynthesis rates, stomatal conductance, and transpiration rates decreased under the drought stress conditions. Drought stress also led to decreased water use efficiency (WUEi). Moreover, leaf proline and soluble carbohydrate contents increased in response to drought stress. While PSII quantum yield reduced under drought stress, it increased under high seed zinc and iron concentrations. It was concluded that enhanced seed zinc and iron concentrations.

Keywords: Chlorophyll fluorescence, Iron, Photosynthesis, Soybean, Zinc.

Introduction:

Optimized use of water is of special importance in agricultural production, particularly in such (semi-) arid regions as Iran, that affects plant growth and development (Tohidloo et al, 2005). Soybean (Glycine max (L.) Merr.) is one of the most important oilseed crops with a relative sensitivity to drought stress (Tarumingkeng and Coto, 2003). The maintenance of the closely related photosynthetic characteristics (i.e., plant growth and yield) is of great significance to the sustainable production of crops. Photosynthesis inhibition leading to reduced crop growth due to drought conditions has been extensively investigated in the literature (Kim et al., 2000; Zlatev and Yorandanov, 2004: Wang et al., 2009). Generally, the first response of plant cell metabolism to root zone water depletion can be seen in its disrupted photosynthetic activities as a result of stomatal and non-stomatal limitations (Flexas et al., 2008). Low intercellular CO₂ concentrations might inhibit not only the RUBISCO and some other photosynthetic proteins (Haupt-Herting and Fok, 2000) but also the activities of ATP synthase enzymes (Noges and Baker, 2000).

Zinc is known to play important roles either as a metal component of enzymes or as a functional, structural, or regulatory cofactor in a large number of enzymes (Grotz and Guerinot, 2006). It also plays an important role in the production of biomass (Cakmak,

*Corresponding Author, Email: Movahhedi1354@yu.ac.ir

2008). Furthermore, zinc takes part in chlorophyll production, pollen function, and fertilization (Pandey et al., 2006). As an activator of Cu-Zn or Mn-SOD, either zinc or manganese is involved in membrane protection against oxidative damages through the detoxification of reactive oxygen species (Marschner, 1995). A high seed Zn content plays very important physiological roles during both seed germination and early seedling growth (Cakmak, 2008). On the other hand, zinc deficiency affects both the photochemical processes of thylakoids and photosynthesis (HajiBoland and Beiramzadhe, 2010), reduces the chlorophyll content, and is likely to cause severe damages to the chloroplast structure (Chen et al., 2007). Studies have reported on reduced stomatal conductance due to zinc deficiency in some plants such as maize (Wang and Jin, 2005) and rice (Hajiboland and Beiramzadhe, 2008).

Iron plays a pivotal role in the different photosynthetic processes of plants. It is, indeed, a key element in CO₂ fixation and biomass production (Briat *et al.*, 2010) while it is also involved in O₂ transport, activation, and detoxification as well as in nitrogen fixation and several photosynthetic reactions. Fe not only contributes to many biological functions and, by altering the ligands to which it is coordinated, has access to a wide range of redox potentials but can also participate in many electron transfer reactions, spanning the standard redox potential (Cairo *et al*, 2002). In order

to perform its multiple tasks, iron must be incorporated in the heme moieties of hemoglobin, myoglobin, and cytochromes; alternatively, it may bind to enzymes in the form of nonheme moieties or Fe–S motifs (such as those associated with several mitochondrial enzymes) (Cairo *et al.*, 2002).

Accumulated proline, known to be a compatible solute, both decreases the cell osmotic potential and helps a sustained water uptake by plants (Hoekstra et al., 2001). Proline reportedly accumulates under a wide range of osmotic stresses (Verbruggen and Hermans, 2008) and its accumulation under drought stress has been well documented in various plant species such as chickpea (Mafakheri, 2010), corn (Serraj and Sinclair, 2002), and peanut (Smith et al., 2002). Accumulation of soluble sugars has also been reported in plants subjected to osmotic stresses despite the fact that the overall plant CO₂ fixation is significantly decreases under osmotic stresses (Murakeozy et al., 2003). Soluble carbohydrates in drought-stressed cells serve such functions as osmotic adjustment, cell protection, carbon reservation, and detoxification (Parvaiz and Satyawati, 2008).

Water use efficiency is defined as the ratio of dry matter accumulation to the amount of water used during the whole growing season, or alternatively, as the ratio of assimilation (A) to transpiration (E) in the short term. This second definition has come to be known as the 'intrinsic water use efficiency' (WUEi) (Hessini et al., 2009). Soil water depletion would cause roots to produce ABA and to translocate it via the xylem to the leaves, thereby decreasing stomatal conductivity (Behera et al., 2002; Liu et al., 2005). Reduced conductivity might in some cases lead to higher levels WUEi. Variations in the parameters Fv/Fm of (maximum quantum yield of PSII) and Fo (initial/minimal fluorescence) are still accepted as reliable diagnostic indicators of the photo-inhibition mechanism. Drought stress has been reported to cause a decline in the maximum quantum yield (Fv/Fm) in wheat (Paknezhad et al., 2007). Adams et al. (2000) maintained that there is an indirect relationship among nutrients (e.g., zinc), chlorophyll synthesis, and electron transfer.

The objectives of this study were to identify the effects of seed zinc and iron concentrations on the photosynthesis, fluorescence parameters, proline content, and total soluble sugars in soybean leaves under drought stress.

Materials and Methods:

A pot experiment was conducted as a split-plot in a randomized complete-block design with three replications. The experiment was carried out in 2012 under greenhouse conditions with a partial shade canopy at Yasouj University, Yasouj, Iran. Drought stress was designated as the main plot with the four levels of well-watered and drought-stressed treatments at the flowering, pod formation, and seed filling stages.

Whole-plot factors were randomly assigned to three blocks. The sub-plot factors consisted of seed nutrient contents with the following eight levels: high seed zinc (78.25 mg kg⁻¹), seed iron (95.57 mg kg⁻¹), seed zinc (77.96 mg kg⁻¹)+iron (94.13 mg kg⁻¹), and the control (Zn 61.02 mg kg⁻¹ and Fe 77.1mg kg⁻¹) as well as seed soaking in distilled water, in 3% zinc sulfate, in 3% iron sulfate, and in 3% zinc+3% iron sulfate solutions. Drought stress was scheduled based on the maximum allowable depletion (MAD) of the available soil water (asw) (i.e., between -0.03 to 1.5 Mpa). Drought stress at each developmental (flowering, pod filling, and seed filling) stage was imposed by irrigation after depletion of 70% of available soil water determined by the weight of each pot (Liu and Stutzel, 2004). The control treatment was irrigated after depletion of only 40% of the available soil water. Seeds with high and low zinc and/or iron contents were obtained from foliar application of zinc and/or iron sulfate in a previous field experiment (Jalil, 2011). Six seeds of the soybean M9 cultivar were sown at a depth of 4 cm in a 25 cm (diameter) \times 35 cm (height) pot containing 7 kg of soil. The plants were subsequently thinned to 3 plants per pot at V_3 stage. The soil had a ratio of 2:1:1 of clay, fine sand, and organic manure, respectively. After five reirrigation events in each drought stress treatment (drought stress at the flowering, pod filling, and seed filling stages), a portable gas exchange system (LCA4, ADC Bioscientific Ltd) was used to measure the including parameters photosynthetic the net photosynthesis rate (Pn), transpiration (E), stomatal conductance to the CO_2 (Gs), and substomatal CO_2 concentration (Ci) using three healthy leaves picked from each pot. Measurements were accomplished in the morning (at 10:00 to 12:00 am) at a temperature of 24 to 35 °C. Chlorophyll fluorescence parameters, including the maximum efficiency of PSII (Fm-Fo/Fm), were measured on three fully expanded young (dark-adapted for 30 min) leaves using a pulse amplitude modulation chlorophyll fluorometer (OS1-FL). The proline content and total carbohydrate sugars of the soybean leaves were measured using the methods described in Paquine and Lechasseure (1979) and Irigoyen and et al. (1992), respectively. The average value of each trait obtained for each pot was calculated and used in the subsequent statistical analysis using the GLM procedure of SAS. Treatment means were compared using Fisher's protected least significant difference (LSD) test (P \leq 0.05).

Results:

Photosynthetic parameters: The effect of drought stress and seed nutrient content interaction was found to be significant ($P \le 0.05$) for all the measured traits, except in the case of Fv/Fm (Table 1). Drought stress severely decreased the value of Pn (Table 2). Seed nutrient content increased Pn at all levels of drought stress. In the non-stress treatment, the highest increase in Pn was observed with the high seed zinc+iron content

Source of								Leaf	Leaf
variation	df	Pn	Е	Ci	Gs	WUEi	FV/FM	proline	Carbohydrate
variation								content	content
Replication	2	0.020 ^{ns}	0.22 ^{ns}	2.65 ^{ns}	0.00004 ^{ns}	0.118ns	0.00007^{ns}	0.227 ^{ns}	213.6240 ^{ns}
Drought stress	3	380.80**	120.34**	79204.02**	0.016**	2.60**	0.0673**	144.197**	1817.3217**
Error a	6	0.248	0.1385	78.72	0.00004	0.050	0.0005	0.6517	92.9888
Element	7	16.81**	7.17**	151353.05**	0.00043**	1.37*	0.0005**	10.54**	1335.1031**
Drought *	21	6 79**	2 21**	7602 20**	0.00046**	1 76**	0.0002 ns	12 107**	1575 2616**
element	21	0.28	5.51	7005.29***	0.0004644	1.20	0.0002	12.107	1373.3010
Error b	56	0.26	0.18	63.2	0.000052	0.074	0.00023	1.52	144.1959
CV		13.49	14.9	6.42	21	21.5	1.99	20.9	0.203

TABLE 1 Analysis of variances for plant characteristics of soybean in the drought stress treatments

*, **, Significant at the 0.05 and 0.01 levels of probability, respectively. ns, not significant.

(116%) compared to the normal seed zinc and iron content (Table 2). The maximum value of Pn in the drought stress treatment at the flowering stage belonged to the ones with a high seed zinc+iron content and a high seed zinc content. In the drought stress treatment at the pod filling stage, the highest increase in Pn was obtained for the high seed zinc content (401%) and for the seed soaking in iron solution (368%) compared to the normal seed zinc and iron content. At the seed filling stage, the maximum value of Pn under drought stress belonged to the treatment containing a high seed zinc content followed by that of seed soaking in the zinc solution and the zinc+iron solution. In this experiment, the lowest value of Pn was observed with the normal seed zinc and iron content treatment.

Values of transpiration rate, E, exhibited a significant decline for all the experimental levels of drought stress. However, the extent of the decline differed among the treatments, with the greatest observed in the drought stress treatments at the seed filling stage and with the seed soaking in iron solution (Table 2). In the non-stress treatment, E was observed to increase in the treatment with a high seed zinc content (75%) and that with a high seed zinc+iron content (66%) compared to the one with a normal seed zinc and iron content (Table 2). Drought stress at the flowering stage and that at the pod filling stage led to increments in E values in leaves with a high seed zinc+iron content (11%) and those from seeds soaked in the iron solution (588%), respectively. Under the drought stress at the seed filling stage, the maximum value of E was observed with both seed soaking in a zinc solution and the high seed zinc and iron content. Zinc and iron concentrations had different effects on the photosynthetic characteristics at each drought stress level. The reduced transpiration and photosynthesis rates were mainly associated with the closure of leaf stomata.

Substomatal CO₂ concentration (Ci) increased in all the drought stress treatments (Table 2). In the non-stress treatments, the lowest and of the highest Ci were observed with seed soaking in iron as well as seed soaking in distilled water and normal seed zinc and iron contents, respectively (Table 2). In the drought stress treatment at the flowering stage, Ci decreased with a high seed zinc content (522%) and with the seed soaking in a zinc solution (394%) compared to those in the normal seed zinc and iron content. In the drought stress treatments at both the pod filling and the seed filling stages, Ci decreased with seed soaking in the zinc and iron solution (101%) and with the high seed zinc content (50%) compared to the normal seed zinc and iron content.

Drought stress severely decreased the value of Gs (Table 2). In the non-stress treatment, Gs increased with the high seed zinc+iron content (107%). In drought stress imposed at the flowering stage, Gs increased with a high seed zinc content (106%) and a high seed zinc+iron content (44%) as well as the seed soaking in a zinc solution (44%) and in a zinc+iron solution (44%) as compared to plants with a normal seed zinc and iron content (Table 2). In the drought stress treatment at the flowering stage, the maximum value of Gs was observed with the high seed zinc content, the high seed zinc + iron content, and the seed soaking in both zinc and zinc + iron solutions. In the drought stress treatment at the pod filling stage and the one at the seed filling stage, maximum values of Gs were observed with both the seed soaking in an iron solution and that in a zinc+iron solution and in distilled water.

WUEi decreased under drought stress (Table 2). In the non-stress treatment, seed soaking in the zinc solution, in the iron solution, and in the zinc+iron solution increased WUEi values by 93%, 92%, and 88%, respectively, compared to those with a normal seed zinc and iron content (Table 2). WUEi increased in the drought stress treatments at the three flowering, pod filling, and seed filling stages with the high seed zinc and iron content (407%), the seed soaking in water (2%), and the seed soaking in iron (887%) compared to the normal seed zinc and iron content.

Drought stress led to reduced maximum quantum yield (Fv/Fm) (Fig. 1). The maximum value of quantum yield was observed in the non-stress treatment (Fig. 1). Maximum values of Fv/Fm were also observed with the high seed zinc content, the high seed zinc + iron content, and the seed soaking in a zinc +iron solution (Fig. 2).

Leaf proline and carbohydrate concentrationDrought stress strongly increased leaf proline concentration (Table 2). In the non-stress treatment, the minimum leaf proline content was observed in the seed soaking in a

TABLE 2 Mean comparisons of the interaction effect s of seed mineral content and drought stress by LSD method									
Drought stress	Element	$\begin{array}{c} Pn \\ (\mu mol \\ m^{-2}s^{-1}) \end{array}$	$(mmol m^{-2}s^{-1})$	(μmol mol ⁻¹)	$(mol m^{-2}s^{-1})$	w θΕι (μmol mol ⁻¹)	w θΕι (µmol mol ⁻¹)	Lear proline content (µmol gr ⁻¹)	Carbohydrate content (mg gr ⁻¹)
e Non stress	Normal seed zinc and iron content	6.21 ^e	5.27 ^c	124.16 ^a	0.053 ^d	1.18 ^b	1.18 ^b	2.31 ^{ab}	50.27 ^{bc}
	High seed zinc content	12.22 ^b	9.25 ^a	7.26d ^e	0.096 ^b	1.32 ^b	1.32 ^b	2.66 ^{ab}	33.1 ^{cd}
	High seed iron content	9.41 ^d	7.11 ^b	31.33 ^b	0.073 ^c	1.34 ^b	1.34 ^b	3.74 ^a	31.8 ^{cd}
	High seed zinc and iron content	13.44 ^a	8.80^{a}	18.5 ^{bcd}	0.110 ^a	1.52 ^b	1.52 ^b	3.49 ^a	34.5 ^{cd}
	Seed soaking in water	5.10^{f}	5.14 ^{cd}	125.96 ^a	0.053 ^d	0.99 ^c	0.99 ^c	2.11 ^{ab}	43.6 ^{bc}
	Seed soaking in zinc solution	12.26 ^b	5.48 ^c	20.80 ^{bc}	0.073 ^c	2.28 ^a	2.28 ^a	2.62 ^{ab}	24.0 ^d
	Seed soaking in iron solution	9.17 ^d	4.06 ^e	5.48 ^e	0.073 ^c	2.27 ^a	2.27 ^a	2.15 ^{ab}	56.7 ^b
	Seed soaking in zinc and iron solution	10.46 ^c	4.69 ^{de}	12.83 ^{cde}	0.056 ^d	2.23 ^a	2.23 ^a	0.67 ^b	76.6 ^a
	Normal seed zinc and iron content	0.48°	1.82 ^{cd}	245.3 ^a	0.016 ^b	0.26^{d}	0.26^{d}	4.68^{f}	26.6 ^c
g stag	High seed zinc content	2.94 ^a	2.59 ^b	39.41 ^f	0.033 ^a	1.12 ^{ab}	1.12 ^{ab}	8.50 ^{cd}	37.9 ^{bc}
Drought stress at flowering	High seed iron content	1.47 ^b	2.07 ^{bc}	79.33 ^d	0.013 ^b	0. 70 ^{bcd}	0. 70 ^{bcd}	6.21 ^{ef}	30.7 ^c
	High seed zinc and iron content	3.75 ^a	2.88 ^a	68.30 ^e	0.023 ^a	1.32 ^a	1.32 ^a	12.32 ^a	56.4 ^{ab}
	Seed soaking in water	1.46 ^b	1.34 ^d	143.43 ^c	0.016 ^b	1.10 ^{ab}	1.10 ^{ab}	7.00^{de}	59.4 ^a
	Seed soaking in zinc solution	1.66 ^b	1.66 ^{cd}	$49/60^{\mathrm{f}}$	0.023 ^a	1.00^{abc}	1.00^{abc}	9.04 ^{bc}	57.7 ^a
	Seed soaking in iron solution	0.92 ^{bc}	1.50 ^{cd}	156 ^c	0.013 ^b	0.62 ^{cd}	0.62 ^{cd}	7.51 ^{cde}	43.8 ^{abc}
	Seed soaking in zinc and iron solution	1.44 ^b	1.61 ^{cd}	178.50 ^b	0.023 ^a	0.88^{abc}	0.88^{abc}	10.66 ^{ab}	40.5 ^{abc}
Drought stress at pod filling stage	Normal seed zinc and iron content	0.6 ^{bc}	0.42 ^d	208.63 ^a	0.020 ^{bc}	1.49 ^a	1.49 ^a	4.60 ^{bc}	38.0 ^b
	High seed zinc content	3.01 ^a	2.24 ^{ab}	159.73 ^d	0.030 ^{ab}	1.33 ^{ab}	1.33 ^{ab}	4.09 ^{bc}	39.9 ^b
	High seed iron content	0.42 ^c	0.49 ^{cd}	191.33 ^b	0.010 ^c	0.87 ^c	0.87 ^c	8.47 ^a	40.8 ^b
	High seed zinc and iron content	0.95 ^{bc}	1.59 ^{bc}	143.33 ^e	0.013 ^c	0.62 ^c	0.62 ^c	10.06 ^a	47.1 ^b
	Seed soaking in water	2.29 ^b	1.54 ^{bc}	176.70 ^c	0.013 ^c	1.52 ^a	1.52 ^a	5.92 ^b	54.1 ^b
	Seed soaking in zinc solution	1.43 ^b	1.52 ^c	125.70 ^f	0.016 ^c	0.94 ^{bc}	0.94 ^{bc}	3.73°	56.8ª
	Seed soaking in iron solution	2.81 ^a	2.89 ^a	111.33 ^g	0.033 ^a	0.95 ^{bc}	0.95 ^{bc}	8.48 ^a	47.9 ^b
	Seed soaking in zinc and iron solution	1.29 ^b	0.92 ^{cd}	103.36 ^g	0.010 ^c	1.38 ^{ab}	1.38 ^{ab}	3.46 ^c	41.1 ^b
ess at seed filling stage	Normal seed zinc and iron content	0.56 ^e	1.33 ^c	151.70 ^d	0.026 ^{ab}	0.41 ^c	0.41 ^c	5.73 ^b	49.5 ^e
	High seed zinc content	3.11 ^a	2.9 ^a	101.20 ^e	0.020 ^b	1.06 ^b	1.06 ^b	8.46 ^a	151.0 ^a
	High seed iron content	2.36 ^{abc}	2.11 ^b	149.70 ^d	0.020 ^b	1.11 ^b	1.11 ^b	4.73 ^b	100.5 ^c
	High seed zinc and iron content	2.09 ^{bc}	1.79 ^{bc}	232 ^b	0.020 ^b	1.17 ^b	1.17 ^b	4.93 ^b	139.9 ^{ab}
	Seed soaking in water	1.64 ^{cd}	1.46 ^{bc}	189.40 ^c	0.033 ^a	1.17 ^b	1.17 ^b	7.44 ^{ab}	90.9 ^{cd}
ıt str	Seed soaking in zinc solution	3.70 ^a	3.57 ^a	154.66 ^d	0.026 ^{ab}	1.03 ^b	1.03 ^b	6.22 ^b	131.2 ^b
1guo.	Seed soaking in iron solution	1.1d ^e	0.28 ^d	291.05 ^a	0.013 ^b	4.05 ^a	4.05 ^a	7.45 ^{ab}	50.9 ^e
Dr	Seed soaking in zinc and iron solution	2.46 ^{ab}	2.13 ^b	147.86 ^d	0.036 ^a	1.15 ^b	1.15 ^b	8.37 ^a	80.1 ^d

TABLE 2 Moon	comparisons o	of the interest	on offect s of s	and minaral cont	tant and draught str	occ by I SD mothe
IADLE 2 WEAT	COMDALISONS O	и ппе ппегасі	on enects of s	eeu mmerai com	LEIIL AIIU UI OUVIIL SLI (ess dy Lodd menn

Means within each column followed by the same latter are not significantly different (α =0.05)

zinc+iron solution. In the drought stress treatment at the flowering stage, leaf proline concentration increased

with the high seed soaking in a zinc+iron solution (163%) compared to the normal seed zinc and iron content. In the drought stress treatment at the pod filling stage, leaf proline concentration increased with the high seed zinc+iron content (119%), seed soaking in an iron solution (84%), and the high seed iron content (83%) compared to the normal seed zinc and iron content. In the drought stress treatment at the seed filling stage, leaf proline concentration increased with the high seed zinc content (57%) and the seed soaking in a zinc+iron solution (46%) compared to the treatment with a normal seed zinc and iron content.

Overall, drought stress increased severely the soluble carbohydrate content of the leaves (Table 2). In the non-stress treatment, at the flowering, pod filling, and seed filling stages, the carbohydrate content increased with seed soaking in a zinc+iron solution (52%), seed soaking in distilled water (123%), seed soaking in a zinc solution (49%), and in those with a high seed zinc content (205%) relative to the normal seed zinc and iron content, respectively (Table 2). Also, in the drought stress treatment at the seed filling stage, an increase was observed in the leaf soluble carbohydrate content.

Discussion:

In this experiment, transpiration rate and stomatal conductance decreased under drought stress conditions at all the reproductive stages of soybean. Zinc and iron different concentrations had effects on the photosynthetic characteristics at each level of the drought stress conditions. The results of this experiment showed that high levels of zinc and zinc+iron contents in seeds led to identical enhancements in stomatal conductance and transpiration rates under drought conditions at the flowering stage. Seed soaking in the iron sulfate solution and a high seed zinc concentration increased stomatal conductance in the drought stress treatment at the pod filling stage. Wang et al. (2009) reported that zinc deficiency in corn reduced the photosynthesic rate, stomatal conductance, and transpiration rate under drought conditions. Zinc may play an important role in increasing stomata openings due to the translocation of large volumes of K⁺ into guard cells. Furthermore, the enhanced Pn of wellwatered soybean plants might be associated with increasing stomatal conductance which, as suggested by Sharma et al. (1995), increases intercellular CO₂ concentrations. On the other hand, it may be claimed that the carbonic anhydrase activity reduced both stomatal conductance and photosynthesis under a zinc deficiency (Hacisalihoglu et al., 2003). It has been shown that adequate zinc nutrition has a protective effect against oxidative reaction caused by reactive oxygen species (Wang and Jin, 2005). Molassiotis et al. (2006) reported that, besides zinc, iron plays an important role in the stomatal opening process. Zinc plays a complex role in various processes and has a close relationship with carbon fixation. In this study, zinc and zinc+iron were observed to have positive effects on the photosynthetic efficiency, even under non-stress conditions. These results are clearly inconsistent with those reported elsewhere (Hajiboland and Amirzadeh, 2010; Zhao and Shen Zhang, 2006; Chuyning *et al.*, 2005; Yin *et al.*, 2006; Felexas *et al.*, 2008).

The present study (Table 2) revealed that Ci increased in the drought stress. Treatment; this might be associated with the low photosynthetic rate. These results suggest that, in addition to the stomatal factors, other non-stomatal factors also affect photosynthestic activities in soybean leaf. The increasing amount of Ci under stress indicates that the metabolic processes involved in carbon fixation are impaired so that cells fail to use efficiently the intracellular carbon. Moradi and Ehsanzadeh (2015) reported that a decline in Pn associated with an increase in Ci has been frequently interpreted as a direct effect of stress on plant photosynthetic capacity.

Zinc and iron nutrients were found to lead to reduced Ci under drought conditions. In general, the lowest Ci was achieved with a high seed zinc concentration, seed soaking in a zinc+iron solution, or seed soaking in a zinc solution in all the three drought stress treatments. This is evidence of the fact that zinc and iron play their protective and structural roles in drought stress conditions to improve upon or maintain photosynthesis. Zinc deficiency has a strong effect on reducing the leaf chlorophyll content; as such, it causes severe damages to the chloroplast structures, especially in drought stress conditions (Chen et al., 2007). Iron is an essential element in the electron transport of photosystem II and is required for chlorophyll synthesis (Ahmadi, 2010). Our results showed that stomatal conductance has a greater effected on Pn than on Ci. Chavez (2004), however, reported that Gs had the greatest effect on Pn under drought stress conditions. Thus, it could be claimed that non-stomatal factors had greater effects on the photosynthesis processes than the stomatal factors did under the drought stress conditions in this experiment. Zinc and iron, especially zinc, were able to enhance photosynthesis under drought stress viatheir effect on stomatal (stomata opening) and non-(detoxification properties, stomatal enabling photosynthetic enzyme activation, etc.) factors.

Another important aspect of drought stress regarding the soybean performance is the intrinsic water use efficiency (WUEi), which is defined as the ratio of photosynthetic rate to transpiration rate (Hessini *et al.*, 2009). The partial stomatal closure under the drought stress conditions led to a greater reduction in transpiration rate than in photosynthesis rate, which resulted in a higher WUEi. The full closure of stomata or damages inflicted on the photosynthesis system would, however, decrease WUEi due to the reduced photosynthesis activities. Zinc and iron nutrition had positive effects on WUEi in soybean plants. This is while the drought stress treatments in the present experiment led to reduced WUEi values in soybean although many authors have reported on improved



Figure 1. Mean comparison of Fv/Fm as affected by drought stress at the different developmental stages of soybean. Means within each column followed by the same latter are not significantly different (α =0.05).



Figure 2. Mean comparison of Fy/Fm as affected by the different seed zinc and iron concentrations in soybean. Means within each column followed by the same latter are not significantly different (α =0.05).

experiment led to reduced WUEi values in soybean although many authors have reported on improved WUEi under water limitation (Hessini et al., 2009; Liu et al., 2005).

Maximum quantum yield (Fv/Fm) reduced under drought stress (Fig 1). The lowest quantum yield was observed in the drought stress treatment at the pod filling stage while its highest value was obtained with high seed zinc+iron concentrations and with seeds soaked in the zinc+iron solution. One reason for the reduced quantum yield or the light inhibition observed is the reduction of leaf chlorophyll content because of a leaf zinc deficiency. Pätsikkä et al. (2002) reported that the total leaf chlorophyll content played an essential role in the imposition of light inhibition in leaves, so that leaves with low levels of chlorophyll were more sensitive to light inhibition. Pn depends on quantum yield so thatquantum yield increases with increasing net photosynthesis rate and the reduction in carbon dioxide

fixation during water stress has been reported to depend on stomatal closure (De Souza et al., 2005).

Drought stress was found to increase proline accumulation in leaves. Maximum proline accumulation observed with the high seed was zinc+iron concentration and with seeds soaked in the zinc+iron solution under drought stress at the flowering and grain filling stages. Proline accumulation helps plants tolerate drought stress (Jaleel et al., 2007; Verbruggen and Hermans, 2008). Askari and Ehsanzadeh (2015) reported that drought stress in fennel (Foeniculum vulgare Mill) led to increased proline and carbohydrate concentrations. The high zinc and iron concentrations were found to have different effects on the leaf soluble sugars in the different drought stress treatments. Seed soaking in a zinc+iron solution in the non-stress treatment led to maximum accumulation of soluble carbohydrates. Generally, seed soaking in each of the elements, compared to the high seed zinc and/or iron

concentration(s), was found more effective in accumulating soluble sugars. Emam Aziz *et al.* (2010) reported that foliar application of the zinc+iron solution led to maximum accumulation of soluble carbohydrates in the Lemongrass plant. Moreover, foliar application of zinc has been reported to increase soluble carbohydrates in safflower under drought stress (Movahhedi Dehnavi, 2004). It was observed in the present study that the higher the soluble carbohydrate content, the less was the Pn and that the reduced soluble carbohydrate activity led to an enhanced photosynthetic rate (Pego *et al.*, 2000). Research has shown that the concentration of such soluble carbohydrates as glucose and fructose

increase the activity of invertase in cells exposed to drought stress (Pinheiro *et al.*, 2001; Trouverie *et al.*, 2003).

Conclusion:

Based on our findings, it might be suggested that increasing seed zinc and/or iron content by foliar application of their solution (s) on the maternal plant or by soaking seeds in the proper solution is a useful method for increasing the photosynthetic parameters in the resulting soybean plants, especially when the plants are exposed to drought stress conditions.

References:

- Adams, M. L., Norvell, W. A., Philpot, W. D., Peverly, J. H. (2000) Spectral detection of micronutrient deficiency in Bragg soybean. Agronomy Journal 92: 261–268.
- Ahmadi, M. (2010) Effect of zinc and nitrogen fertilizer rate on yield and yield components of oilseed Rape (*Brassica napus* L.). World Applied Sciences Journal 10: 298-303.
- Askari, E and Ehsanzadeh, P. (2015) Drought stress mitigation by foliar of salicylic acid and their interactive effects on physiological characteristics of fennel (*Foeniculum vulgare* Mill.) genotypes. Acta Physiologiae Plant 37:4.
- Emam Aziz, E., Ezz El-Din, A., Omer, E. A. (2010) Influence of zinc and iron on plant growth and chemical constituents of *Cymbopogon citrates* L. grown in newly reclaimed land. International Journal of Academic Research 2: 278-283.
- Behera, R. K., Misha, P. C., Choudhury, N. K. (2002) High irradiance and water stress induce alterations in pigment composition and chloroplast activities of primary wheat leaves. Journal of Plant Physiology 159: 967–973.
- Briat, J. F., Duc, C., Ravet, K. and Gaymard F. (2010) Ferreting and iron storage in plant. Biophysica Acta General Subjects 1800: 608-814.
- Bukhov, N. G. (2004) Dynamic light regulation of photosynthesis (a review). Russian Journal of Plant Physiology 51: 742-753.
- Cairo, G., Recalcati, S., Pietrangelo, A. and Minotti, G. (2002) The iron regulatory proteins: targets and modulators of free radical reactions and oxidative damage. Free Radical Biology & Medicine Journal 32:1237–1243.
- Chaves, M. M. and Oliveira, M. M. (2004) Mechanisms underlying plant resilience to water deficits: prospects for water-saving agriculture. Experimental Botany 55: 2365–2384.
- Cakmak, I. (2008) Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? Plant and Soil 302: 1-17.
- Chen, W., Yang, X., He, Z., Feng Y. and Hu F. (2007) Differential changing photosynthetic capacity, 77K chlorophyll fluorescence in chloroplast ultra-structure between Zinc-efficient and Zinc inefficient rice genotypes (*Oryza sativa* L.) under low Zinc stress. Journal of Plant Physiology 132: 89–101.
- Chunying, Y., Youhong, P., Runguo, Z., Yaping, Z. and Chunyang, L. (2005) Adaptive responses of *Populus kangdingensis* to drought stress. Journal of Plant Physiology 123: 445-451.
- De Souza, C. R., Maroco, J. P., dos Santos, T. P., Rodrigues, M. L., Lopes, C., Pereira, J. S., Chaves, M. M. (2005) Control of stomatal aperture and carbon uptake by deficit irrigation in two grapevine cultivars. Agriculture, Ecosystems & Environment Journal 106: 261-274.
- Flexas, J., Bota, J., Galmés, J., Medrano, H. and Ribas-Carb, M. (2008) Keeping a positive carbon balance under adverse conditions: responses of photosynthesis and respiration to water stress. Physiologica Plantarum 127: 343-352.
- Hacisalihoglu, G., Hart, J. J., Wang, Y., Cakmak, I. and Kochian, L. V. (2003) Zinc efficiency is correlated with enhanced expression and activity of Cu/Zn superoxide dismutase and carbonic anhydrase in wheat. Journal of Plant Physiology 131: 595-602.
- Hajiboland R. and Beiramzadeh, N. (2010) Growth, photosynthesis and antioxidant defense system in Zn-deficient *red cabbage* plants. Plant, Soil and Environment Journal 56: 209–217.
- Hajiboland, R. and Beiramzadeh, N. (2008) Growth, gas exchange and function of antioxidant deironnse system in two contrasting rice genotypes under Zinc and Iron deficiency and hypoxia. Acta Biologica Szegediensis 52: 283–294.
- Haupt-Herting, S. and Fok, H. P. (2000) Exchange of oxygen and its role in energy dissipation during drought stress in tomato plant. Plant Physiology 110: 489-495.
- Hessini, K., PabloMartiniz, J., Gandour, M., Albouchi, A., Soltani, A. and Abdelly, C. (2009) Effect of water stress on growth, somatic adjustment, cell wall elasticity and water- use efficiency in *Spartina alterniflora*. Environmental and Experimental Botany 67: 312-319.

- Hoekstra, F. A., Golovina, E. A. and Buitink, J. (2001) Mechanisms of plant desiccation tolerance. Trends in Plant Science 6: 431–438.
- Irigoyen, J. J., Emerich, D. W. and Sanchez-Diaz, M. (1992) Water stress induced changes in concentration of praline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants. Physiologia plantarum 84: 55-60.
- Jaleel, C. A., Gopi, R., Manivannan, P., Kishorekumar, A., Sankar, B., Panneerselvam, R. (2006) Paclobutrazol influences vegetative growth and floral characteristics of *Carthamus roseus* (L.). Indian Journal of Applied and Pure Biology 21: 369- 372.
- Jalil, M. (2011) Effect of foliar application of zinc and iron on quantity and quality yield of Soybean (*Glycine max* L.) under drought stress. M. Sc, Yasouj University, Yasouj, Iran. (Language in Persian).
- KaviKishor, P. B. (2005) Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. Current Science 88: 424–438.
- Kim, J. Y., Mahe, A., Brangeon, J. and Prioul, J. L. (2000) A maize vacuolar invertase, IVR2, is induced by water stress. Organ/tissue specificity and diurnal modulation of expression. Journal of Plant Physiology 124: 71–84.
- Liu, F., Andersen, M. N., Jacobsen, S. E. and Jensena, C. R. (2005) Stomatal control and water use efficiency of soybean (*Glycine max* L.) during progressive soil drying. Environmental and Experimental Botany 54: 33-40.
- Liu F., Jensena, C.R. and Andersen, M. N. (2004) Drought stresses effect on carbohydrate concentration in soybean leave sand pods during early reproductive development: its implication in altering pod set. Field Crops Research 86:1–13.
- Liu F. and Stutzel, H. (2004) Biomass partitioning, specific leaf area and water use efficiency of vegetable amaranth (*Amaranthus* spp.) in response to drought stress. Scientia Horticulturae 102: 15-27.
- Mafakheri, A., Siosemardeh, A., Bahramnejad, B., Struik, P. C. and Sohrabi, Y. (2010) Effect of drought stress on yield, proline and chlorophyll contents in three chickpea cultivars. Australian Journal of Crop Science 4:580-585.
- Maschner H. (1995): Mineral nutrition of higher plant. Second edition, academic press Inc London, PP. 891.
- Maxwell, K. and Johnson, G. N. (2000) Chlorophyll fluorescence. A practical guide. Experimental Botany 51: 659-668.
- Molassiotis, A., Tanou, G., Diamantidis, G., Patakas, A. and Terios I. (2006) Effects of 4-month Iron deficiency exposure on Iron reduction mechanism, photosynthetic gas exchange, chlorophyll fluorescence and antioxidant defense in two peach rootstocks differing in Iron deficiency tolerance. Plant Physiology 163: 176–185.
- Moradi, L and Ehsanzadeh, P. (2015) Effects of Cd on photosynthesis and growth of safflower (*Carhamus tinctorius* L.) genotypes. Photosynthetica 53: 506-518.
- Movahhedi Dehnavi, M. (2004) Effect of foliar application of micronutrients (zinc and manganese) on the quantitative and qualitative yield of different autumn safflower cultivars under drought stress in Isfahan. Ph.D. Thesis, Tarbiat Modarres University, Tehran, Iran.
- Murakeozy, E. P., Nagy, Z., Duhaze, C., Bouchereau, A. and Tuba, Z. (2003) Seasonal changes in the levels of compatible osmolytes in three halophytic species of inland saline vegetation in Hungary. Plant Physiology 160: 395–401.
- Noges, S., Baker, N. R. (2000) Effect of drought on Photosynthesis in Mediterranean plant grown under Enhanced UV-B radiation. Journal of Experimental Botany 51: 1309-1317.
- Paknejad, F., Nasr, M. R., Tohidi Moghadam, H., Zahedi, H. and Jami Alahmad, M. (2007) Effects of drought stress on chlorophyll fluorescence parameters chlorophyll content and grain yield of wheat cultivars. Biological Sciences 7: 841-847.
- Paquine, R. and Lechasseure, P. (1979) Observation surune metode dosage la libre dance les de planttes. Canadian Journal of Botany 57: 1851-1859.
- Parvaiz, A., Satyawati, S. (2008) Salt stress and phyto-biochemical responses of plants a review. Soil and Environment 54: 89–99.
- Patsikka, E., Kairavuo, M., Sersen, F., Aro, M. E. and Tyystjarvi, E. (2002) Excess copper predisposes photosystem II to photoinhibition in vivo by outcompeting iron and causing decrease in leaf chlorophyll. Plant Physiology 129: 1359–1367.
- Pego, J. V., Kortstee, C., Huijser, S. C. and Smeekens, M. (2000) Photosynthesis, sugars and the regulation of gene expression. Experimental Botany 51: 407–416.
- Pandey, N., Pathak, G. C. and Sharma, C. P (2006) Zinc is critically required for pollen function and fertilisation in lentil. Journal of Trace Elements in Medicine and Biology 20: 89-96.
- Pereira, E. W., Siquera, L. D., Martinez, C. A. and Puiatti, M. (2004) Gas exchange and chlorophyll fluorescence in four citrus root stocks under aluminum stress. Plant Physiology 157: 513-520.
- Pinheiro, C., Chaves, M. M. and Ricardo, C. P. (2001) Alterations in carbon and nitrogen metabolism induced by water deficit in the stems and leave of Lupinus albus L. Journal of Experimental Botany 52: 1063–1070.
- Serraj, R. and Sinclair, T. R. (2002) Osmolyte accumulation: Can it really help increase crop yield under drought conditions? Plant, Cell and Environment 25: 333 -341.
- Sharma, P. N., Tripathi, A. and Bisht, S. S. (1995) Zinc requirement for stomatal opening in caulifower. Plant Physiology 107: 751–756.

- Smith, B. N., Girija, C., Swamy, P. M. (2002) Interactive effects of sodium chloride and calcium chloride on the accumulation of proline and glycine betaine in peanut (*Arachis hypogaea* L.). Environmental and Experimental Botany 47: 1-10.
- Tarumingkeng, R. C. and Coto Z. (2003) Effects of drought stress on growth and yield of soybean. Science Philosophy PPs 702, Term paper, Graduate School, Borgor Agricultural University (*Institut Ppertanian* Bogor), December.
- Tohidloo, G. h., Sadeghian, S. Y., Kashani, A., Gohari, J., Taleghani, D. F. & Hamdi, F. (2005) Study on water use efficiency, some agronomical and physiological characteristics on the three lines of sugar beet in well watered and stress conditions. Journal of Agronomy and Crop Science 191: 279-301.
- Trouverie, J., The'venot, C., Rocher, J. P., Sotta, B. and Prioul, J. L. (2003) The role of absisic acid in the response of a specific vacular invertase to water stress in the adult maize leaf. Journal of Experimental Botany 54: 2177–2186.
- Verbruggen, N., Hermans, C. (2008) Proline accumulation in plants: a review. Journal of Amino Acids 35:753-759
- Wang, H. and Jin, J. Y. (2005) Photosynthetic rate, chlorophyll fluorescence parameters, and lipid peroxidation of maize leaves as affected by zinc deficiency. Photosynthetica 43: 591-596.
- Wang, H., Liu, R. L. and Jin, J. Y. (2009) Effect s of zinc and soil moisture on photosynthetic rate and chlorophyll fluorescence parameters of maize. Biologia plantarum 53: 191-194.
- Yin, C. Y., Berninger, F. and Li, C. Y. (2006) Photosynthetic responses of *Populus przewalski* subjected to drought stress. Photosynthetica 44: 62-68.
- Zhao, F. and Shen Zhang, B. (2006) Physiological foundation for the difference of long term water use efficiency among *Populus deltoids* clones. Acta Ecologica Scientica 26: 2079-2086.
- Zlatev, Z., Yordanov, I. (2004) Effects of soil drought on photosynthesis and chlorophyll fluorescence in common bean plants. Bulgarian Journal of Plant Physiology 30: 3-18.